

Used as directed at 300 ppm concentration (one 0.56g tablet dissolved in one quart of warm water) in waters up to 500 ppm of hardness calculated as CaCO₃, this product is effective against the following pathogenic organisms in 60 seconds:

Community Associated Methicillin Resistant *Staphylococcus aureus* (CA-MRSA), *Escherichia coli*, *Escherichia coli* 0157:H7, HIV-1 (AIDS Virus), *Listeria monocytogenes*, *Staphylococcus aureus*, *Shigella dysenteriae*, and *Yersinia enterocolitica*.

SANI-SPRAY sanitizing solutions of 300 ppm concentration may be used on food processing equipment, utensils and other food contact articles and surfaces as specified under 40 CFR 180.940 of the US Environmental Protection Agency.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS and DOMESTIC ANIMALS

WARNING: Tablet dust causes substantial but temporary eye injury. Do not get dust in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Tablets may be harmful if swallowed.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN: Rinse immediately with water for 15-20 minutes. If irritation persists, call poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by poison control center doctor. Do not give anything by mouth to an unconscious person. Have the product label or container with you when calling a poison control center or doctor, or going for treatment.

ESL09052012

SANI-SPRAY

MULTI-PURPOSE SANITIZING TABLETS



USED AS DIRECTED, EACH 0.56g TABLET PRODUCES ONE QUART OF NO RINSE FOOD CONTACT SURFACE SANITIZER SPRAY SOLUTION

Kills HIV-1 (AIDS Virus) and Community Associated Methicillin Resistant *Staphylococcus aureus* (CA-MRSA) when used as directed for sanitizing.

ACTIVE INGREDIENT: (Quaternary) Alkyl (C14 95%, C12 3%, C16 2%)
Dimethyl benzyl ammonium chloride dihydrate 50%
OTHER INGREDIENTS 50%
TOTAL 100%

**KEEP OUT OF THE REACH OF CHILDREN
WARNING**

See left panel for additional precautionary statements.
For Institutional and Commercial use only.

EPA REG. No. 1561-11 EPA EST. No. 1561-VA-1

**200
TABLETS**

Manufactured by
EDWARDS-COUNCILOR CO., INC. NET WT
1427 Baker Road, Virginia Beach, VA 23455 **4 OZ**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Sani-Spray Tablets is a no-rinse food contact surface sanitizer.

DIRECTIONS FOR SANITIZING WITH HAND TRIGGER SPRAY BOTTLE

FOR SANITIZING FOOD PROCESSING EQUIPMENT, DAIRY EQUIPMENT, SINKS, COUNTERTOPS, TABLES, REFRIGERATED STORAGE AND DISPLAY EQUIPMENT AND OTHER HARD NON-POROUS FOOD CONTACT ARTICLES AND SURFACES.

1. Wash and rinse all articles and surfaces thoroughly.
2. Apply a solution of 300 ppm concentration by dissolving one tablet (0.56g) in a clean one quart size (32 oz.) hand trigger spray bottle filled with warm water, allowing several minutes for tablet to dissolve thoroughly before using. Spray surfaces to be sanitized with hand trigger spray bottle and allow surfaces to remain wet for at least one minute followed by adequate draining and air drying. Do not rinse or wipe.

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS

Personal Protection: The specific barrier protection items to be used when handling items soiled with blood or body fluids are disposable gloves, gowns, masks and eye coverings.

Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

Contact Time: One minute.

Disposal of Infectious Materials: Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

STORAGE AND DISPOSAL. Do not contaminate water, food or feed by storage and disposal. Store in original container in areas inaccessible to persons unfamiliar with its use. Nonrefillable container. Do not reuse or refill this container. Triple rinse promptly after emptying. Offer empty container for recycling if available or discard in trash. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

